

Onondaga County Health Department

Division of Environmental Health  
421 Montgomery Street  
Syracuse, New York 13202

# **Incinerator Monitoring Program**

## **2010 Summary Report**

June 1, 2011

**Submitted To: Cynthia B. Morrow, M.D., M.P.H.**  
Commissioner of Health

**Submitted By: Kevin L. Zimmerman**  
Director, Division of Environmental Health

**Summary Statement:**

In the monitoring conducted to date, no relationship has been established between the operation of the incinerator and any significant increased levels of constituents in the environment.

## **Introduction**

The Onondaga County Health Department initiated an incinerator monitoring program in 1994, the year prior to the Waste to Energy (WTE) facility being placed into operation. In 2003, the monitoring program for air, soil and ash was reevaluated, and a more effective and efficient program was developed and implemented starting in 2004. As an alternative to offsite air monitoring, direct interaction was established with the Onondaga County Resource Recovery Agency (OCCRA) and the New York State Department of Environmental Conservation (DEC) in providing stack monitoring results and improved assurance on reporting of adverse events and equipment failures. This allows for evaluation of short-term changes in the incinerator emissions, an effective alternative to the previous limited scope offsite air monitoring conducted over a nine year period.

Long-term deposition impacts continue to be evaluated by soil and ash monitoring. All soil samples are analyzed for metals twice a year. Several changes related to organics testing have been implemented based on the low levels detected in the monitoring conducted to date, and the fact that there is no evidence of a trend or levels associated with health risks. In 2009, half of the soil sampling sites were analyzed for organics during the spring collection. The remaining sites were tested during 2010. The monitoring program has the flexibility of testing a site again in the following year should an elevated level of any organic constituent be detected. The four soil ash route sites have been eliminated from the program. To date these sites have not shown any elevation of metals or organics indicating that ash transport in covered vehicles is not a significant environmental or health concern. Ash, directly from the incinerator continues to be analyzed for metals twice a year and organics once a year. Under present contracts, organic analysis is performed by Axy's Analytical Services, LTD, and metal analysis is performed by Life Science Laboratories, Inc. The collection of soil is performed by Onondaga County Health Department, Division of Environmental Health staff, while collection of the ash is the responsibility of Covanta Energy System under New York State Department of Environmental Conservation protocols.

## **Air Monitoring**

During 2010, the department interacted directly with OCCRA and DEC in review of the stack monitoring results and reporting of adverse events and equipment failures by the facility operator, Covanta Energy. The department also reviewed both the monitoring conducted at the stack on a continuous basis and reported quarterly to DEC, as well as the annual stack test that is performed by an independent contractor. At no time did the monitoring indicate constituents above levels of health concern. The annual stack test incorporates an extensive list of analytes that include metals and organics. All of the analytes were well below permit limits.

## **Soil and Ash Testing for Organics**

Soil from seven routine soil sites collected in the spring of 2010 was analyzed for dioxins/furans (PCDD/PCDF), polychlorinated biphenyls (PCB's), and polycyclic aromatic hydrocarbons (PAH's). Ash, also collected in the spring of 2010, was analyzed for the same constituents.

Organic sample results are compared to published background data and U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry (ATSDR) Toxicological Profiles, EPA Preliminary Remediation Goals, and NYSDEC Soil Cleanup Objectives. In general, little change in levels of these compounds have been observed from background through the present organic screening period. The levels of organics in the ash were similar to reports for ash identified by other investigators and reported in published literature.

Each form of dioxin/furan has associated with it a toxic equivalency factor that is used to calculate a total toxic equivalency (TEQ) for each sample. Attachment A shows the historical dioxin/furan TEQ values for routine soil sites and ash samples. All levels remain well below the ATSDR and EPA action levels and there is no indication of a trend. For ash, dioxin/furan total TEQ remain fairly consistent. Ash is not homogeneous and an inconsistent result occurs occasionally. The results are similar to those reported by other investigators.

Attachment B shows the historical PCB values for routine soil sites and ash samples. All levels remain below the ATSDR and EPA action levels and there is no indication of a trend. One site, Syracuse University, had a PCB level that was elevated compared to the last sampling period. This contamination may have been caused by past activities on this property. The site will be carefully monitored during the next sampling period.

## **Soil and Ash Testing for Metals**

Soil from the fourteen soil sites and ash were analyzed for ten different metals twice during the year (Spring and Fall). The metal results are issued in two different reports, one for soils and one for ash.

Metal results are compared to background levels, published national averages for urban areas and a statewide rural soil survey. Soil and ash are not homogeneous and can contain materials that can account for an occasional inconsistent result. Hence, a single elevated or depressed value will not be assumed to be indicative of a change at a specific site. Rather, the pattern of values for that specific element must demonstrate a statistically significant difference, which may be indicative of a real environmental change. In general, the metal results for 2010 fall within the expected range of values for urban areas and demonstrate no significant variation from background levels.

Attachment C shows the historical levels for the ten metals at the routine soil sites. Due to the volume of data, the mean (average) of all routine sites and all routine control sites for each year is presented. The complete report includes all of the data for each site. Attachment C-1 provides data on New York State Department of Environmental

Conservation Soil Cleanup Objectives, a New York State rural soil survey, and USEPA soil screening levels for metals in residential soil.

Attachment D shows the historical levels of the ten metals in ash.

### **Summary and Conclusions**

In general, the organic and metal results for this monitoring period are within the expected range for urban environments and are below any levels associated with health risk. Any fluctuations in sample results appear to be a reflection of the low levels detected, expected variation as a result of sample collection, preparation, and laboratory procedures, or possible variable levels due to past activities at a site. All levels remain below those associated with health concerns. The results should be viewed in the context of an ongoing program of environmental monitoring performed by the Onondaga County Health Department as a part of its overall Incinerator Monitoring Program. In the monitoring conducted to date, no relationship has been established between the operation of the incinerator and any significant increased levels of constituents in the environment.

The following are the detailed Incinerator Monitoring Program reports that have been issued on the 2010 soil and ash testing:


- 2010 Screening Summary for Organic Constituents
- 2010 Soil Metals Analysis Summary
- 2010 Ash Characterization Summary

Copies of these reports are available upon request.

The following abbreviations may be used in this report:

As	Arsenic.
ATSDR	Agency for Toxic Substances and Disease Registry
Be	Beryllium.
Cd	Cadmium.
CES	Certified Environmental Services.
Cr	Chromium.
CV	Coefficient of Variation.
ELS	Environmental Laboratory Services.
Hg	Mercury.
LD	Limit of Detection.
ND	None Detected.
ug/g	micrograms per gram.
Ni	Nickel.
OCCF	Onondaga County Correctional Facility.
OCHD	Onondaga County Health Department.
PAH	Polyaromatic Hydrocarbon
PCB	Polychlorinated Biphenyls
PCDD/PCDF	Polychlorinated Dibenzo-p-Dioxins/Dibenzofurans
Pb	Lead.
pg/g	picograms per gram
PPM	parts per million.
SD	Standard Deviation.
Se	Selenium.
SHFD	Sentinel Heights Fire Department
V	Vanadium.
WTE	Waste to Energy Facility.
Zn	Zinc.
~	approximately.
<	Less than.
>	Greater than.
NA	Not applicable.
NS	Not sampled.

# OCHD ROUTINE SOIL MONITORING SITES



Attachment  
A

Dioxin/Furan TEQ Soil Results through Year 2010 (pg/g dry weight)

Routine Soil Sites

Site	Year											
	1994	1999	2000	2001	2002	2004	2005	2006	2007	2008	2009	2010
Clark Reservation	1.8	1.2	2.27	1.42	1.23	2.03	1.90	1.76	1.73	1.26	***	1.64
Jamesville Beach	0.6	0.5	1.09	0.82	0.70	0.71	0.97	0.86	0.93	0.77	***	0.52
OCCF	0.79	2.2	1.88	1.47	1.26	1.38	5.54	1.52	1.94	1331.72@	1.72	***
DOT @ Jagaith	2		1.5	1.64	3.41	2.41	3.78	3.38	1.73	39.90@	2.62	***
Dutch Hill *	0.77		1.41	1.16	1.40	1.03	1.26	1.02	1.02	0.64	***	0.73
Erie - Poolsbrook *	1.39		1.5	1.14	1.86	**	**	**	**	**	**	**
Nottingham	0.51		0.78	0.79	0.80	0.70	0.94	0.85	0.84	0.74	0.76	***
SHFD	12		8.02	9.89	9.72	7.02	8.09	6.27	7.20	10.74	***	7.12
Sevier Rd	1.8		2.07	2.58	2.56	**	**	**	**	**	**	**
Beaver Lake *			0.51	0.53	0.85	0.70	0.72	0.64	0.69	0.65	0.38	***
Ch. 3 Towers			3.36	3.88	3.35	9.66	7.79	7.69	5.39	2.44	3.72	***
Gen. Crushed Stone			2.77	1.98	2.13	**	**	**	**	**	**	**
Highland Forest			1.18	1.24	0.96	**	**	**	**	**	**	**
JD High School			1.32	1.29	1.12	1.10	1.48	1.16	1.06	1.28	***	1.13
Nob Hill			0.93	0.91	0.90	6.83	1.01	1.00	1.07	1.05	***	0.78
Pratts Falls			0.91	0.98	0.77	0.87	0.98	0.83	0.94	1.17	0.82	***
Southwood			0.6	1.14	1.01	1.08	1.05	0.97	1.09	1.01	0.80	***
Syracuse University			3.11	6.97	9.47	13.89	3.14	3.66	12.96	0.67	***	2.45

\* Denotes Control Sites  
 \*\* Site no longer sampled due to program re-evaluation  
 \*\*\* Site not sampled this year. Sites are sampled every other year.  
 @ A single elevated value will not be assumed to be indicative of a change at a specific site, rather a pattern of values must demonstrate a statistically significant difference.

Combined Ash

Site	Year											
	1999-Spring	1999-Fall	2000-Fall	2001-Fall	2002-Fall	2004-Spring	2005-Spring	2006-Spring	2007-Spring	2008-Spring	2009-Spring	2010-Spring
Day 1 and 2	256	153	109	123	177	72	191	246	250	243	168	200
Day 3, 4, and 5	242	205	154	137	220	445	142	148	276	240	126	172

Note: For reference purposes, the ATSDR investigation level for Dioxin/Furan TEQ is 50 pg/g and the EPA clean up level is 1,000 pg/g.



Attachment  
B

## PCB Results through Year 2010 (pg/g dry weight)

## Routine Soil Sites

Site	2000	2001	2002	2004	2005	2006	2007	2008	2009	2010
Clark Reservation	6010	2360	3150	2780	3610	2770	4110	2840	***	2860
Jamesville Beach	1260	644	683	703	1110	781	1220	1610	***	589
OCCF	3080	5230	2000	2310	6940	3120	6320	2190	2810	***
DOT @ Jaquith	16100	15400	45100	9220	67100	49100	18000	14200	34700	***
Dutch Hill *	2210	1170	1400	1200	1380	1140	1450	1340	***	1060
Erie - Poolstrook *	2620	1400	2020	**	**	**	**	**	**	**
Nottingham	2140	2280	3610	1640	7380	2850	3050	2110	4200	***
SHFD	3080	2970	1760	1900	2730	1610	2510	1730	***	2240
Sevier Rd	1870	1600	2250	**	**	**	**	**	**	**
Beaver Lake *	1970	1210	5250	2650	1420	1360	1360	1370	2450	***
Ch. 3 Towers	3360	2310	2490	1620	1830	1730	2220	1400	1510	***
General Crushed Stone	9430	3160	5450	**	**	**	**	**	**	**
Highland Forest	2120	1210	1270	**	**	**	**	**	**	**
JD High School	3580	1780	1732	1810	2640	1780	1720	2720	***	1750
Nob Hill	3500	2480	2500	3440	2810	2970	2830	2950	***	2510
Pratts Falls	1890	1840	1440	1620	1650	1220	1450	2050	1230	***
Southwood	2240	2160	1150	1480	1470	1470	2750	1640	1640	***
Syracuse University	10700	114000	11000	9510	6940	11400	10900	1170	***	78600

\* Denotes Control Sites

\*\* Site no longer sampled due to program re-evaluation

\*\*\* Site not sampled this year. Sites are sampled every other year.

## Combined Ash


Site	2000-Fall	2001-Fall	2002-Fall	2004-Spring	2005-Spring	2006-Spring	2007-Spring	2008-Spring	2009-Spring	2010-Spring
Day 1 and 2	79000	22000	13600	7850	2470	5770	3080	23000	3100	5930
Day 3, 4, and 5	4700	7020	6580	38000	33000	57000	3060	5550	51900	8840

PCB results prior to 2000 were all less than detection limits. Starting in 2000 detection limits were lowered so that usable concentrations were available.


Note: For reference purposes, the ATSDR indicates that typical mean PCB concentrations in background soil are less than 100,000 pg/g

Attachment  
C


VII.A. Comparison of Annual Mean Values  
Routine and Routine Control Sites



## VII.B. Comparison of Annual Mean Values Routine and Control Sites




### VII.C. Comparison of Annual Mean Values Routine and Control Sites






### VII.D. Comparison of Annual Mean Values Routine and Control Sites




VIII.E. Comparison of Annual Mean Values  
Routine and Control Sites







### VII.F. Comparison of Annual Mean Values Routine and Control Sites



VII.G. Comparison of Annual Mean Values  
Routine and Control Sites




### VII.H. Comparison of Annual Mean Values Routine and Control Sites






VIII.1. Comparison of Annual Mean Values  
Routine and Control Sites



VII.J. Comparison of Annual Mean Values  
 Routine and Routine Control Sites



# **Attachment C-1**

Metal	NYS SCO's for restricted use residential (ppm)	Rural Soil Survey (ppm)	USEPA Soil Screening levels for residential (ppm)
Arsenic	16 (0.21)	16	0.39
Beryllium	14	1.2	160
Cadmium	2.5 (0.86)	2.5	70
Chromium	36	30	280
Lead	400	133	400
Mercury	0.81	0.3	6.7
Nickel	140	29.5	1600
Selenium	36	4	390
Vanadium	NA	38	390
Zinc	2,200	180	23,000

New York State Department of Environmental Conservation Soil Cleanup Objectives. The Health Based SCO's were calculated considering all exposure pathways: ingestion, inhalation, dermal, carcinogenic (1 in a million cancer risk), and non-carcinogenic (using risk reference doses). The final health based SCO is based on the most conservative pathway calculation. In some cases the SCO has been modified to match background if the rural background levels for NYS are above the calculated SCO (the health based SCO is in parenthesis). Restricted use means no livestock or animal product consumption.


NYS Statewide Rural Surface Soil Survey (2005)-determined concentration ranges for 170 commonly assessed analytes in discrete surface soil samples collected at randomly selected rural NYS properties.

USEPA Soil Screening Levels for residential—Values were calculated based on the ingestion-dermal exposure pathway for residential soils. These screening levels are not action levels or clean up levels, they are a tool for further evaluation.


# Attachment D



VII.A. Mean Values Ash Data  
Dry Weight



VII.B. Mean Values Ash Data  
Dry Weight



VII.C. Mean Values Ash Data  
Dry Weight

